

Issue on 10.1016/j.mssp.2023.107881

The SEM equipment was mis-identified on this article [1]. The authors stated on the Method section that the SEM images were obtained with a SU8010 model (Hitachi), however, the images show that they were obtained with a Regulus 8100 model.

10.1016/j.mssp.2023.107881

5 GH Foundation

and compositions of CQDs/CO/PFO photocatalysts. A SU8010 fieldemission scanning electron microscopy (FE-SEM) and JEM 2100F transmission electron microscopy (TEM) were used to observe the sur-




Fig. 4. FE-SEM photographs of CQDs/CO/PFO photocatalysts. (a) 3 mL-CQDs/CO/PFO, (b) 5 mL-CQDs/CO/PFO, (c) 7 mL-CQDs/CO/PFO, and (d) 9 mL-CQDs/CO/PFO, and (d) 9 mL-CQDs/CO/PFO, (e) 7 mL-CQDs/CO/PFO,

The 5GH Team wants to address that misidentifying the SEM equipment does not necessarily suggest that the authors got involved in misconduct, but the authors from India (Navdeep Dhaliwal, Satbir S. Sehgal), and Saudi (Mohd Ubaidullah, Asad Syed) are questionable.

Article Information

Title: CQDs drives CeO2/PbFe12O19 photocatalysts for oxytetracycline hydrochloride removal by photoinduction and magnetic recovery

Authors: Xiangyu Chen, Shifa Wang (王仕发, 三峡学院), Yujia Jin, Maoyuan Li, Huajing Gao, Hua Yang, Leiming Fang, Mohd Ubaidullah, Navdeep Dhaliwal, Satbir S. Sehgal, Asad Syed, Dengfeng Li

Journal: Materials Science in Semiconductor Processing

DOI: 10.1016/j.mssp.2023.107881

[1] 10.1016/j.mssp.2023.107881

This article is licensed to the 5GH Foundation under a CC BY-NC-ND 4.0 International License.